

Information provision and network externalities: the impact of genomic testing on the dairy industry

Victor Funes-Leal¹, Jared Hutchins¹

¹*Department of Agricultural and Consumer Economics, University of Illinois*

Link to the latest version:

<https://victorefunes.github.io/files/job-market-paper.pdf>

Abstract

We estimate the impact of a new testing technology: genomic selection, on the market for dairy cattle genetics in the United States. The genomic selection allows cattle breeders to test new cattle as soon as they are born for the presence of desirable, productive traits and more than double their pace of genetic improvement in milk production. At the same time, the rate of inbreeding, the extent to which dairy bulls are related to one another, greatly increased after the introduction of genomic testing. Using a data set of all bulls marketed in the US from 2000 to 2020, we examine whether inbreeding rates increased due to increased competition for popular dairy cattle families in the face of greater market competition. We use a difference-in-difference and matching approach and find that inbreeding increased greatly in already popular families after the introduction of genomic testing relative to less popular families with similar production and health traits. Our estimation shows that the increased inbreeding rate of American bulls caused a loss of between 3.6 to 6.7 billion dollars to the entire industry from 2011 to 2019.

1 Introduction

The United States is the world's most important supplier of cattle genetics. In 2022, the United States alone exported \$295 million of bovine genetics, representing 47% of the world's exports in value terms. This means that the continued health and quality of the US cattle genetics gene pool is not only important to US livestock industries but also to livestock industries all across the world.

In terms of increasing output per cow, the US dairy cattle genetics industry is seemingly a success story. For about six decades, milk production per cow has consistently increased in the US herd, and about half of this growth has been attributed to improvements in cattle genetics. Dairy breeders improve milk yield per cow by first identifying which bulls have the most productive female offspring and then continuing to breed off of these lines. In previous years, the pace of this genetic improvement was limited by biological factors. Determining whether a bull had good milk production ability took at least three years since the bull had to have offspring born from their use on dairy farms and those offspring would need to take two years to grow to production age.

In the past ten years, an innovation in testing technology has revolutionized the industry: genomic testing. Introduced in 2008, genomic testing allows breeders to link each animal's genes and productive outcomes such as milk yield and fat yield. The practical upshot of this is that determining a dairy bull's quality can now be done as soon as it is born instead of in three years. There were two important impacts of genomic testing in the market for dairy bulls. First, the pace of genetic improvement more than doubled in dairy cattle since breeders could find productive bulls at much younger ages and breed new generations even quicker. Second, dairy companies began more than doubling the amount of bulls for sale on the market. Because genomic testing allows faster genetic discovery, some scientists theorized that genomic testing would lead to a decrease in inbreeding, measured as the degree of relatedness of an animal to all others, as more diverse sets of genetics are discovered and incorporated into the gene pool.

The introduction of genomic testing, however, has coincided with a less desirable outcome: an increase in inbreeding. Inbreeding rate is a measure of the relatedness of a

dairy bull to the rest of the dairy cattle population. Inbreeding is an example of a negative externality in the cattle genetics industry since higher inbreeding rates expose the industry to more disease risk and depressed yields in future generations, costs not borne directly by the breeder in the short run. Since dairy breeders continually breed from each other's genetic stock, it is also the case that the breeding decisions of one breeder will affect all other breeders through the shared public resource that is the genetic pool available in the US. In one case, breeding excessively from a famous dairy bull "Pawnee Farm Arlinda Chief" led to more than 500,000 spontaneous abortions and nearly 420 million dollars in losses because of a deleterious mutation in a gene that was passed to his offspring (Adams et al., 2016). Given the US role in the global cattle genetics market, damaging the diversity of the dairy cattle gene pool could cause damage to livestock markets all around the world.

In this paper, we test whether the introduction of genomic testing caused an increase in competition for already popular dairy bull genetic lines and, thus, an increase in the inbreeding rate. Our analysis is complicated because popular dairy bull lines are usually superior to other lines in terms of production and health traits. To get a clean comparison, we construct a counterfactual line that is comparable to popular lines in terms of productivity using propensity score matching. Using a difference-in-difference approach, we then examine inbreeding rates in popular families relative to our counterfactual line before and after the introduction of genomic testing in 2010. If genomic testing increased competition for quality lines independent of previous popularity, we expect that inbreeding rates for these our two groups will be the same both before and after its introduction.

We find that, after the introduction of genomic testing, popular dairy bull lines became more inbred than our counterfactual. Relative to our control group, the inbreeding rate of the most popular lines increased 1-2.8 percentage points after 2010. Before genomic testing, there was no statistically significant difference in inbreeding rates across our groups. Using an estimate of the costs of inbreeding from the Holstein breed association, we estimate that this increase in inbreeding led to \$3.6-6.7 billion in losses to the industry from 2011-2019. Since dairy breeders are incentivized to inbreed for short-run gains even at the expense of their genetic pool, this indicates a role for regulation to penalize the industry for inbreeding in the short run to prevent damages to the industry in the long

run.

This article fills a critical gap in the literature; there are no other papers that analyze inbreeding as a **network externality** (Liebowitz and Margolis, 1994), (Katz and Shapiro, 1985), where companies enjoy the benefits (faster genetic gains) and suffer the costs (decreased fertility caused by the increased frequency of deleterious mutations) from changes in the size of their associated network. In this case, the pedigree (relatedness) is the force that binds all the nodes in the network, where two animals are part of the same network (line) if they descend from the same ancestor. Similarly, all articles that analyze the market for cattle genetics, such as Kerr (1984), Melton et al. (1994), Richards and Jeffrey (1996) or Schroeder et al. (1992) study cattle as a closed system and their objective was to determine what are the implicit prices of each genetic trait using hedonic methods. These early papers then focus on the decisions taken by individual breeders or dairy farmers without paying attention to their broader impact.

The paper is structured as follows: Section 2 presents the relevant background details and a detailed causal mechanism to explain why introducing genomic selection led to higher inbreeding rates in subsequent generations of dairy bulls. Section 3 explains the data sources and shows how some descriptive statistics can help us delineate the relationship between our observed variables. Section 4 outlines the identification strategy and methodological framework used to identify the causal effect of genomic selection on inbreeding rates. Finally, in section 7, we explain the results and discuss their implications for the future of the American dairy industry.

2 Conceptual Framework

Selective breeding has been practiced since prehistoric times; however, the discovery of inheritance mechanisms and genetics in the 19th and 20th centuries allowed breeders to identify the link between desired traits in animals and their inheritance mechanisms (Lush, 1937). From an economic standpoint, an animal breeder supplies a product (animal genetics) with an inseparable bundle of traits embodied in a single animal. To produce such an animal, the breeder combines a set of inputs that can be split into two groups:

a series of inputs exogenously supplied by the producer (labor, feed, capital, veterinary services) and a series of inputs embodied in the animal (traits.) Those inputs are combined on a production function subject to the prices of exogenous inputs (observed) and genetic inputs, whose prices are implicit but can be estimated based on the price paid for a bull's genetics using the hedonic approach (Schroeder et al., 1992, Sy et al., 1997).

Melton et al. (1994, 1979) add an additional element: breeders are paid based on the (positive) deviation of the trait levels of their bulls relative to the population average. Breeders compete to maximize profits; to do so, they will attempt to release bulls into the market with the highest possible trait values. Dairy farmers demand genetics to improve their productivity in the next period by choosing the bulls with the best characteristics. However, the traits that constitute an animal are marketed as a "bundle" of traits embedded in each animal. Dairy farmers are only interested in the productivity of their cows because their profits depend on it; they are not directly interested in how inbred their cows are; therefore, inbreeding rates cannot be priced directly as an additional cost to dairy farmers.

Inbreeding is the mating of individuals more closely related to each other than the average for the population (Bourdon, 2000, p. 337). Inbreeding has two consequences: first, it increases the likelihood that an animal's offspring will perform similarly to its ancestors; for example, if a particularly tall bull is mated with a related cow, its offspring will likely be taller than average because the genes that carry such trait are likely to be present in both parents. Second, it favors the expression of deleterious recessive alleles; since an animal has two copies of a recessive gene from each parent, it is more likely to express any such traits than an outbred animal.




Figure 1: Inbreeding Rates and Total performance Index

Figure 1 shows the close link between the **Total Performance Index (TPI)**, a weighted average of a series of genetic traits, and the average inbreeding rate. Genomic selection led to an immediate jump in the productivity index but to a lagged response in inbreeding rates because of the gestation length and time to reach maturity that originate the cattle cycle (Rosen et al., 1994), breeding decisions in year t , leading to a birth of a new calf in $t + 1$, that will be released into the market in year $t + 2$.

The increment in the rate of change in hereditary traits caused by genomic selection can be decomposed into a set of components: heritability, selection intensity, and generation interval (Kerr, 1984). Equation 1 is called the “breeder’s equation”:

$$\text{genetic improvement} = \frac{\text{heritability} \times \text{selection intensity} \times \text{trait variance}}{\text{generation interval}} \quad (1)$$

Heritability is a population parameter specific to each trait that measures the strength of the relationship between the genetic traits of parents and their offspring, and so is the

standard deviation of the observed trait values. **Selection intensity** is the differential in average trait values between selected animals and the general population at a given time. Selection intensity measures how “choosy” breeders are in deciding which individuals are selected; more intensity of selection means that only the best animals (according to some criterion) are selected to have offspring. If animals were randomly selected, those magnitudes would not differ. The generation interval is the time required to replace one generation with the next: the shorter the interval, the faster the genetic change. Genetic improvement is driven by two forces: selection intensity and generation interval. An increase in selectivity or a decrease in the generation interval increases genetic gains per generation.

Before 2009, it took one to five years for a bull to father a large enough number of daughters to test them for the presence of genetic traits such as milk, protein, or fat yields. With genomic testing, an animal can be tested as soon as it is born for the presence of specific traits in its genome. Consequently, the generation interval, the average age of parents when offspring are born (Wiggans and Carrillo, 2022), has decreased from around five and a half years to less than two years, close to the biological minimum of one year.




Figure 2: Average age at first proof for Holstein bulls
 Source: NAAB

The most important consequence of a shorter generation interval is an increase in genetic progress. Figure 2 also shows that the number of daughter-proven bulls has steadily decreased, and the wider confidence intervals for the average age reflect that. A key element of our analysis is the existence of what we call “supersires”, high-prestige bulls who became the founders of lines (families). The more genetics companies breed the same lines, the more newborn bulls will be related to a specific ancestor. We define a bull as a supersire in terms of the size of its offspring; since we have records of all male descendants from each sire that have been released on the dairy genetic market, we can then identify which bulls have had the largest offspring; more precisely, a bull born between 1997 and 2007 is a supersire if the size of his offspring is on the 95th percentile of the offspring distribution.

We test the hypothesis that genomic selection caused higher inbreeding rates by constructing a counterfactual line of bulls that possessed similar traits but were less popular

at the introduction of genomics. We identify their descendants and respective inbreeding rates; they increased disproportionately more for the descendants of supersires. A likely explanation for this process is that genetics companies did not invest in the most productive bulls but instead in the most famous lines in an attempt to gain market share. The more companies breed into those lines, the higher the inbreeding rate of the next generation of bulls. This leads to a deterioration of the expected traits of future generations of bulls and, thus, to significant losses for the dairy industry in terms of reduced productivity.

3 Data Description

Improved genetics via the selection of bulls (sires) and cows (dams) is a crucial driver for productivity improvements in the dairy industry. Milk production in the United States has tripled in the last forty years, and over half of this increase is solely due to genetics (<https://uscpcb.com/impact/>). The United States is also the world's largest exporter of bovine genetics, with a share of 46.4% of the total value of exports in 2019. Every improvement in genetics in the United States will be quickly transmitted to the rest of the world dairy industry.




Figure 3: Share of trade Value of Bovine Semen exports by country (2019)

Source: UN Comtrade database

Traditional selection techniques relied on information from pedigrees and observed trait values to assess an animal’s fitness. Genomic selection uses data from an animal’s genome to accurately predict breeding values, particularly in cases where traits are expressed only in females. In such case, it was necessary to measure the traits of a large number of daughters from the same bull; for that to happen, these cows must have reached the age of sexual maturity (one year); hence, the entire process could take three to four years until enough information was gathered.

Instead, genomic selection uses the bull’s DNA to detect gene combinations significantly different from the expected frequency if they were randomly associated. This process detects the presence of certain genetic traits that are highly correlated with specific gene combinations. It can be undertaken as soon as the calf is born, so by the time the animal has reached sexual maturity (one year old), it can be marketed immediately.

Figure 5 shows that genomic selection increased the number of animals released into the market since there is intense competition for supplying the most productive bulls; the relative value of a line is now higher than ever. Since dairy farmers have a more extensive

choice set of bulls to buy genetics from, lines act similar to "brands," as they reduce the quality uncertainty from a product. Finally, this "loyalty" to established lines leads to breeders supplying more bulls from such families than from lesser-known lines that have similar or better traits.

Brand loyalty arises in this context due to the nature of animal genetics, dairy farmers demand a particular set of genetic traits related to the productivity and health of their cows, but they cannot be sold separately; they have to be acquired as a "bundle" at a single price (Ladd and Martin, 1976, Melton et al., 1994) that is a linear function of the entire set of genetic traits, which makes a comparison between any two or more animals extremely difficult. This implies that there are very high search costs to farmers when deciding which animal to buy genetics from, Stigler (1961) and Farley (1964) argue that it is reasonable for an economic agent to seek further information on a single purchase as long as the expected reduction in price per unit as a result of additional search times the quantity purchased is equal to the marginal cost from searching for lower prices (or higher qualities.)



Figure 4: Total number of new bulls released in the market **Source:** NAAB

Genomic selection increased the number of new bulls released into the market, as shown in Figure 4. Consequently, search costs for dairy farmers increased as well, thus promoting greater brand loyalty for "superstar" lines, that is, bulls descended from a notorious ancestor (e.g., bulls that fathered many daughters). The ultimate consequence of this process is an increase in inbreeding for the entire population of dairy bulls; the more related bulls are, the more likely their offspring is to preserve their genetic traits. Higher inbreeding levels lead to a higher likelihood of observing **inbreeding depression**, a decrease in the performance of inbred animals due to lower fertility, higher incidence of rare diseases, and reduced longevity.

The National Dairy Herd Improvement Program (NDHIP) collects, manages, and analyzes data from American dairy cattle (Hutchins and Hueth, 2023). The information it collects assesses the industry's productivity and estimates the paternal contribution to these traits. The raw data collected from farms is used to estimate a statistical model to predict the genetic contribution of an animal as a sire (or dam) after controlling for

environmental variables such as herd, age, pedigree, and season effects. The predicted values for each trait from this model are commonly called the **predicted transmitting ability** (PTA) of a trait.

Animal breeders and genetics companies compete to offer an animal with the most significant possible improvement over the cohort’s average. As a consequence, firms will try to produce bulls with the highest productivity traits; similarly, this will lead to higher rates of inbreeding since the best way to ensure high productivity is through linebreeding, that is, by building a pedigree that has a relatively stable degree of relationship relative to a supersire. We call “large” companies those with the most significant average number of bulls across the entire 2000-2020 period in the North American genetics market. Similarly, a firm is called “preexisting” if it participated in the bull genetics markets before 2009; otherwise, they are considered “new.”




Figure 5: Number of genotyped bulls by firm size
Source: NAAB

Genomic testing allows genetic companies to detect suitable bulls earlier; thus, they

can release those animals into the market as soon as they reach sexual maturity. Figure 5 shows the total number of new bulls released in the market by year and type of firm.

In the cattle genetics industry, genomic testing has been adopted remarkably fast. Figure 6 shows how fast genomic testing was adopted in our sample. The high starting costs for genotyping help explain this since large companies were the ones that could initially afford it. Still, more companies have genotyped their calves since genomic selection was released. Larger firms genotyped a more significant percentage of their newborn calves in 2010, and the share has grown steadily, while smaller firms followed them at a slower rate.




Figure 6: Share of each status of bull by year

Starting in 2010, large firms started genotyping their bulls, and consequently, the number of new bulls increased threefold in that year, when all bulls aged 1 to 4 were released simultaneously. The staggered adoption of genomic selection has further consolidated the market and led to more competition for superstar lines. However, the more inbred selected animals are, by definition, also increases the inbreeding of an average animal in

any cohort, thus making inbreeding depression more likely within a herd. Figure 5 shows the total number of new bulls released in the market by year and type of firm. We call “large” companies with the most significant share of the North American genetics market. Similarly, a firm is called “preexisting” if it participated in the market of bull genetics before 2009; otherwise, they are considered “new.”

4 Empirical Framework

We want to test the hypothesis that the introduction of genomic selection led to higher inbreeding rates because of increased demand for well-known pedigrees fueled by an increment in the productivity of dairy cattle. We want to know the causal effects of introducing this technology; ideally, we need a treatment group that was affected by genomic selection and a treatment group that was not.

We must identify families whose founders had similar trait levels as superstars but had less “momentum”; that is, a smaller offspring. This is a comparable group because breeders could have invested in them but did not because they were not demanded as highly as superstars. We can track these lines over time. If their inbreeding rates did not differ substantially from those of superstars, we could argue that genomic selection allowed breeders to identify new, less-inbred bulls independent of their ancestry.

Figure 6 shows that the pace of adoption was very high from the beginning; this makes it unlikely to use non-genomic proven bulls as a counterfactual, mainly because we don’t know why breeders choose not to genotype certain bulls after 2009. Similarly, we have very few foreign-born bulls to use as a control group; we also don’t know the selection criteria to import genetics from those animals. Both alternative control groups suffer from an additional issue: they will have a significantly smaller offspring, thus leaving us with little statistical power to detect any relevant treatment effect. To overcome these issues, we use propensity score matching (Smith and Todd, 2005) to define a control group that resembles the treatment in all observable characteristics (except for our outcome: inbreeding rates.)

We define a **line** as the set of direct (male) descendants from a particular bull, which,

in turn, can be described in terms of his number of male descendants; a bull is a superstar if his number of sons is in the 95th percentile of the distribution of offspring. Breeders are under high competitive pressure to supply bulls with the highest PTAs. However, the best way to do so is by linebreeding a family of bulls so that the relative distance to an ancestor remains constant across generations. Steyn et al. (2022) and Steyn et al. (2023) show that younger sires capture a more significant share of present-day population genetic variation; hence, genotypes of older sires can be discarded from (genomic) evaluations. As generations begin to accumulate, the contribution of a particular sire decreases (on average) exponentially, so we can disregard the impact of distant ancestors.

In the sample, we include all animals whose sires were born between 1998 and 2004, and we call those sires the “founders” of each line. Our definition depends critically on when the founders of such lines were born; if we choose an earlier date, we cannot find a statistically significant effect due to the high relatedness of all bulls. The reason behind this is that bulls are advertised only with their sire and grandsire; any ancestry beyond that is not taken into account by dairy farmers. If we choose a later date, both treatment and control groups will have little difference in inbreeding rates because there is not enough time for the control group lines to develop. Figure 12 in the appendix summarizes the sample selection mechanism; the treatment and control groups are comprised of the descendants of two groups: supersires (treatment) and matched sires (control).

After identifying the sires, we can recursively build their pedigrees by matching bulls with their sires until we find the latest animal in the sample. We want to measure the causal effect of introducing genomic testing in 2009 on the average inbreeding in extant lines; the first problem we must tackle is the nature of a valid comparison group. The introduction of genomic testing caused animal breeders to supply more animals from well-known lines, that is, descended from superstar sires; our control group must be the lines descended from other non-superstar sires.

First, we need to match the founders of such lines with other sires that did not father as many sons but had comparable genetic traits. Second, we find all descendants from both groups and compare the evolution of their inbreeding rates through time. The idea behind this procedure is that when those bulls were in the market, any reasonably similar sire

could have been chosen but was not, and thus, can be used to construct a counterfactual line.

For every family, we must construct a counterfactual family with new bulls being born every year, each with an inbreeding rate that we can average across cohorts. Then, the **treatment group** are all lines of animals comprised of direct descendants of superstar sires. Similarly, the **control group** includes descendants from comparable animals matched to superstar bulls by their genetic trait levels.

Our next objective is to identify the counterfactual lines by matching superstar sires with alternative lines. To do so, we use a propensity score matching estimator, let $p(X) = \Pr(T_j = 1|X)$ where T is a dummy equal to 1 if sire j 's offspring is on the 95th percentile of the distribution and X_j be a vector comprised of the following:

1. Predicted Transmitting Ability for production traits (milk, fat, and protein), health traits (mastitis, metritis, productive life), and reproductive traits (daughter pregnancy rate, effective calving, gestation length).
2. Dummies for the absence of certain recessive genetic disorders (Holstein Foundation, 2021) such as *brachyspina*, cholesterol deficiency, and mule foot.
3. Dummies for the presence of genes associated with specific qualitative traits (polled, red coat color, complex vertebral malformation).
4. Dummies for the presence of haplotypes (specific sequences of DNA at different locations in the chromosome) that affect fertility (Holstein Foundation, 2018) when present in homozygous form (likely to be expressed when two carrier animals are mated).

We then estimate the propensity score (Cameron and Trivedi, 2005, p. 873), the probability of being a superstar given the level of PTAs and the presence of the entire set of variables listed above. The validity of our method relies on a series of assumptions (Heckman et al., 1997):

Assumption 1 (Stable Unit Treatment Value Assumption): $(inb^{treatment}, inb^{control}) \perp\!\!\!\perp T/X$.

The average inbreeding rates of any two lines do not vary with the treatment assigned to other lines. For each unit, there are no different forms of each treatment level (Imbens and Rubin, 2015), conditional on a set of covariates. This assumption implies that conditional on \mathbf{X} , the distribution of inbreeding rates for both groups will be the same for all founders, but it must differ for their offspring. Figure 13, in the Appendix, shows the extent to which the inbreeding rates distribution of our matched sample resembles that of the treatment group. Unlike unmatched sires, their inbreeding distribution has a similar mean and variance conditional on the set of genetic traits.

Assumption 2: $0 < \Pr(T_j = 1 | \mathbf{X}) < 1$ for all \mathbf{X}

This assumption implies that we can define a propensity score for every value in the treatment group. Assumptions 1 and 2 together are the “strong ignorability” assumption in Rosenbaum and Rubin (1983), and they are necessary in conjunction to identify causal parameters. The set of matching variables above exhausts the entire set of possible selection criteria for breeders; no unobserved variable can influence mating decisions. Figure 14 shows the degree of overlapping between superstars and alternative sires; since there is a considerable difference in the size of the groups, superstars make up only a tiny fraction of bulls; however, the overlapping is good in the range 0.05 – 0.50, but there exist a few superstars that had little close matches in the sample.

A central part of our argument is that treatment and control groups should have similar inbreeding levels before genomic selection. To test that assumption, in Table 1, we calculate the difference in averages of treatment and control groups for inbreeding rates and a few selected traits. There is no statistically significant difference for most traits between treatment and control groups, so our matching method ensures that the difference between both groups is due to the “prestige” of superstar lines because the bulls in the treatment group had a larger offspring than those in the control group. Each year these bulls will have a new set of descendants being born, each with its inbreeding rate and genetic traits; if our hypothesis holds, then the inbreeding rates of the descendants of supersires should be significantly higher than those of non-superstar descendants.

We then estimate the differential impact of genomic selection (our treatment) on the two types of lines, those from supersires and those from alternative sires. Model (2)

includes leads and lags of the treatment to check to which degree a pre-treatment balance exists in both groups. This is not a test for parallel trends on our differences-in-differences specification (Cunningham, 2021), but it helps us ascertain how comparable both groups are. Table 1 shows that there is no significant pre-treatment difference in both groups and that coefficients from 2010-2011 are not statistically significant either, which coincides with the values from Figure 1 where there is a significant increment in inbreeding rates starting in 2012.

We observed the average inbreeding rates of the treatment and matched control groups before and after introducing genomic selection. If we divide the 2005-2017 period into two groups, pre-intervention (2005-2009) and post-intervention (2010-2018), denoted as 1 and 2 and let T_j denotes whether a bull from line j belongs to treatment ($T_j = 1$) or control ($T_j = 0$) groups, then:

$$inb_{jt} = T_j inb_{jt}^{treat} + (1 - T_j) inb_{jt}^{control}$$

Where $j = 1, \dots, J$ are the lines, each comprised of n_j bulls such that $i = 1, \dots, n_j$. Our causal estimand of interest is the Average Treatment on the Treated (ATT) in the post-intervention period:

$$ATT_{post} = E \left[inb_{post}^{treat} - inb_{post}^{control} \middle| T_j = 1 \right]$$

This quantity measures the average causal effect on the treated lines in the post-treatment period (Roth et al., 2023). To correctly identify causal effects, a series of assumptions must hold (Sun and Abraham, 2021):

Assumption 3 (Parallel trends):

$$E \left[inb_{post}^{control} - inb_{pre}^{control} \middle| T_j = 1 \right] = E \left[inb_{post}^{control} - inb_{pre}^{control} \middle| T_j = 0 \right]$$

We assume that the average inbreeding rates for both groups would have evolved in parallel without introducing genomic selection. This assumption is not testable because we cannot observe untreated inbreeding rates after 2009, but it constrains our model

specification to a Two-Way Fixed Effects specification we call Model (1):

$$inb_{ikt} = treat_i + (treat_i * post_t)\beta + post_t + \alpha_k + \varepsilon_{ijt} \quad (\text{Model (1)})$$

Where inb_{ikt} is the inbreeding rate of bull i born from a line that belongs to either the treatment or control group born in year t , supplied by firm k , β_t is a year of birth fixed effect, and α_j is a firm fixed effect.

Assumption 4 (No anticipatory effects): $inb_{pre}^{control} = inb_{pre}^{treat}$

This assumption ensures that both groups were comparable before the treatment; inbreeding rates of superstar lines did not increase before the introduction of genomic selection; we use matching methods to ensure that this condition holds.

Under Assumptions 3 and 4, we have that:

$$ATT_{post} = E \left[inb_{post}^{treat} - inb_{post}^{control} \mid T_j = 1 \right] - E \left[inb_{pre}^{treat} - inb_{pre}^{control} \mid T_j = 0 \right] \quad (2)$$

The ATT is identified as the difference-in-differences of observed outcomes in the pre and post-intervention periods. All elements in this equation are observed and, therefore, estimable. Suppose superstars had superior traits to our matched control group. In that case, this condition is violated, and we would not be able to parse out the causal effect of inbreeding from the effect of the bull's genetic superiority. Assumptions 3 and 4 ensure that the OLS estimates $\hat{\beta}$ from Model (2) are consistent and asymptotically valid (Roth et al., 2023).

Assumption 5: Let $W_j = (inb_{j,pre}, inb_{j,post}, T_j)'$ be a vector of outcomes and treatment status for unit i . Then, we observe a sample of N independent and identically distributed random draws $W_i \sim F_W(w)$ for a distribution F satisfying parallel trends.

Under Assumptions 3, 4, and 5: $\sqrt{N}(\hat{\beta} - ATT_{post}) \xrightarrow{D} N(0, \sigma^2)$, then we can consistently estimate the variance σ^2 clustering at the line level (Bertrand et al., 2004) asymptotically ($N \rightarrow \infty$ and T fixed).

We can also decompose the variations in treatment time since every year, a certain number of bulls will be born from lines in both groups, each with its inbreeding rate that can be averaged across treatment and control groups. Models 2-4 show the complete

specification with leads and lags of the treated group. These models decompose the estimated ATT relative to the treatment average by birth year. Sun and Abraham (2021) call this specification “Dynamic specification.” However, there is no staggered treatment since genomic selection affects all lines in the same year, but inbreeding rates evolve independently after the treatment.

This assumption states that the treatment status of an individual from any given line is not affected by the treatment status of others. In our setting, the treatment (genomic selection) affects all individuals, so there is no selection into treatment. To measure the effects of genomic selection on inbreeding, we estimate the differences-in-differences model:

$$inb_{ikt} = treat_i + treat_i * \sum_{\substack{t=2005 \\ t \neq 2009}}^{2019} \beta_t \mathbf{I}(yob = t) + \alpha_t + \alpha_k + \varepsilon_{ijt} \quad (\text{Model 2})$$

$$inb_{ikt} = treat_i + treat_i * \sum_{\substack{t=2005 \\ t \neq 2009}}^{2019} \beta_t \mathbf{I}(yob = t) + \alpha_t + \alpha_k + \mathbf{X}_{it} \gamma + \varepsilon_{ijt} \quad (\text{Model 3})$$

$$inb_{ikt} = treat_i + treat_i * \sum_{\substack{t=2005 \\ t \neq 2009}}^{2019} \beta_t \mathbf{I}(yob = t) + \alpha_t + \alpha_k + \mathbf{X}_{it} \gamma + (\mathbf{X}_{it} * post_t) \rho + \varepsilon_{ijt} \quad (\text{Model 4})$$

Model (2) is a specification that does not contain any covariates (PTAs), and it decomposes the ATT on inbreeding rates by year of birth cohort. Model (3) incorporates PTAs (\mathbf{X}_{it}) and firm-level fixed effects, and Model (4) includes interactions of PTAs and the post-treatment dummy. We include the last specifications to account for trait-level trends. We also know that the treatment affects both inbreeding rates and PTAs, but we do not have a structural model for their impact (Lewbel, 2019), so we include them in a non-linear fashion as both levels and interactions to reduce the uncertainty about the functional form of our controls.

5 Results

Figure 7 shows treatment and control groups' average inbreeding rates and confidence intervals across years. Genomic selection did not increase inbreeding rates immediately; both curves began to diverge substantially after 2011. Inbreeding rates, in turn, started growing two years later, and both curves diverged significantly from 2012 onwards.




Figure 7: Average inbreeding rates in treatment and control groups

This plot closely matches the evolution of the population average inbreeding rates plotted in Figure 1. This behavior is consistent with our hypothesis that breeders prefer well-known over new lines. Our findings coincide with results reported by Guinan et al. (2023), who also look at the time trends of productivity and reproduction traits in US dairy cattle. The authors also document a decrease in the generation interval and an increase in inbreeding rates of Holstein and Jersey cattle but do not consider the latter an issue because the expected inbreeding rates of future offspring adjust genetic trait values.

Table 2 shows the results of three models; model (1) is the analog to Figure 7, including

only the interactions between treatment and year of birth. Model (2) splits the treatment effect by year of birth averages, Model (3) includes the complete set of genetic traits and firm-level fixed effects, and Model (4) is the model with traits and a post-treatment dummy. In all specifications, standard errors are clustered at the line level to better deal with the inconsistency generated by the autocorrelation of inbreeding rates between different generations (Bertrand et al., 2004).

Model 1 is the simple differences-in-differences estimation where $post_t$ is a dummy equal to 1 if the bull was born after 2009. The results show that bulls descending from supersires increased their inbreeding rates by 1.46 percent more than those who descend from less popular bulls, on average. Since the impact of genomic testing on inbreeding rates only became noticeable after 2011, the coefficient underestimates the ATT of genomic selection. Model 2 shows that the treatment-year of birth interactions are positive and significant from 2012 onwards, with the 2018 effect being the highest.

Models 3 and 4 in Table 2 show that there is a positive and significant incremental variation for the treated versus control groups in the years 2012-2018; however, the addition of controls in either levels or interactions attenuates the coefficients for 2014 and 2015 to the point of reducing the significance of interactions between treatment and year of birth. Still, in all instances, the coefficients are positive. Regression models show a positive and increasing inbreeding rate in the treatment group comprised of all descendants from prestige sires relative to the control group of descendants of comparable non-superstar sires.

The last period in the sample is 2018 instead of 2020 because we cannot make inferences after 2018 due to a reduction in the number of animals in the control sample; the number of comparable descendants is too small for us to consider them a valid comparison group. This drastic reduction is a consequence of lower fertility rates in the control group; since those sires fathered fewer sons, their sons will beget even fewer grandsons, and after a few years, this group collapses.

Table 2 also includes the results for the F-test for joint significance of all pre-treatment differences in trends for the interaction of treatment and year of birth. The p-values show no significance in two of the three specifications and a 5% significance for the pre-treatment

coefficients of Model 4. This is not a test for parallel trends since the assumption relies on unobservable counterfactuals, letting $\hat{\beta}_k$ be the coefficient for any of the post-treatment periods, then:

$$E[\beta_k] = ATT_k + \left(E \left[inb_{ik}^{control} - inb_{i0}^{control} \middle| T_i = 1 \right] - E \left[inb_{ik}^{control} - inb_{i0}^{control} \middle| T_i = 0 \right] \right)$$

The parallel trends assumption implies that the term in parentheses is equal to zero, then the coefficients correctly capture the ATT for any post-treatment period $k = t - 2009 > 0$. On the other hand, its pre-treatment analog is identified as:

$$E[\beta_{(-k)}] = E \left[inb_{i(-k)}^{control} - inb_{i0}^{control} \middle| T_i = 1 \right] - E \left[inb_{i(-k)}^{control} - inb_{i0}^{control} \middle| T_i = 0 \right]$$

We have shown that these coefficients are not significantly different from zero for all pre-treatment periods, individually and jointly. It is usually considered evidence in favor of the parallel trends assumption. However, this is not necessarily the case since this test has low power against a linear or quadratic pre-trend alternative (Roth, 2022). In our case, we can dismiss this concern; since our matching method ensures the fulfillment of the condition for the founders, the entire effect must be due to their sons and grandsons.




Figure 8: Coefficient plots for different specifications

Figure 8 shows the coefficients from the interaction of year fixed effects and treatment status to assess the impact of inbreeding per year correctly; every year after 2012, the inbreeding rate of the treatment group is between 1 and 2.8 percent points higher than the control group. The coefficient values decrease slightly after controlling for their genetic traits but remain statistically significant. The figure also shows that all pre-treatment coefficients are not significant; we interpret this as evidence in favor of the reasonability of the research design. Figure 8 uses the *sup-t* uniform confidence intervals for the pre-treatment periods of Montiel-Olea and Plagborg-Møller (2019) as suggested by Freyaldenhoven et al. (2019) instead of the standard pointwise confidence. These confidence intervals contain the actual path of the coefficients 95% of the time and can better help understand which types of pre-trends are consistent with the data.

So far, we have discussed our results regarding inbreeding rates only, but we can attach a monetary value to increased inbreeding rates. The American Holstein Breeders Association estimates that every additional inbreeding percent over the average is worth 23 US dollars per animal. The total cost depends on what we consider the average inbreeding; we can use the control group rates as a counterfactual and calculate the cost

from increased inbreeding of the treatment group, multiplied by the total stock of Holstein cattle; in that instance, we estimate a total cost from increased inbreeding of 3.6 billion dollars for the period 2011-2019. Alternatively, by industry standards, the maximum recommended inbreeding rate is 6.25%¹, so we can calculate how much is the cost from inbreeding levels greater than such value, we estimate the total cost for the same period as 6.7 billion dollars for the entire dairy industry.

6 Robustness Checks




Figure 9: Coefficient plots for quarter of birth effects

Since we have the exact date of birth of every animal in the sample, we can group them by a quarter of birth as well; Figure 9 shows the results from Models 2-4 but replacing year for a quarter of birth interactions with the treatment. The signs and magnitudes of coefficients do not differ substantially from the original formulation. The increase in

¹<https://www.agproud.com/articles/36810-the-impact-of-genomics-on-rapid-increase-of-inbreeding-of-holsteins>

coefficient values is monotonic and statistically significant since 2012, but having a smaller sample size per quarter makes confidence intervals wider.

This article’s results depend heavily on a somewhat arbitrary definition of a “superstar” bull in terms of the size of his line. What would happen if we changed that from the 95th to another quantile of the line size distribution? Figure 10 shows the coefficient plots for different definitions of “superstar”.




Figure 10: Robustness test: changes in the definition of “superstar”

If we change the cutoff point from the 95th to the 99th percentile, the sign, and magnitude of our coefficients do not change much. Still, the confidence intervals widen significantly because our treatment group comprises such specific animals that the matching algorithm cannot find suitable matches. As a consequence, the significance of the difference-in-differences coefficients decreases significantly. Going in the other direction, we can find marginally better matches for the superstar group. However, the control group ended abruptly in 2015 because most of the original group’s control lines are now included in the treatment, making the control group short-lived. We conclude that our results depend critically on a set of lines that allow for a good comparison when included in the control group; otherwise, the analysis yields insignificant results.




Figure 11: Coefficients from model (2) per different definitions of founders' date of birth

Finally, Figure 11 shows coefficient plots by different definitions of the founders' date of birth. We defined founders as all bulls born between 1997 and 2005, but such a definition is also arbitrary. If we adopt different cutoffs for the period where the founders were born from 1995/2000 to 2003/2008, we have $6 \times 6 = 36$ distinct combinations for starting and end date. The positive and significant effect of genomic selection on inbreeding rates will arise if the starting date is from 1995 onwards. Still, it rarely persists for lines whose founder was born after 2008. The rationale behind such results is that all lines born after 2008 will have their founders active during the introduction of genomic selection, thus making the effect entirely endogenous.

7 Discussion

This article investigates the consequences of genomic selection on the American dairy industry. On the one hand, this new technology significantly improved productivity, but on the other hand, it increased the average inbreeding rate of all bulls in the market. Such increment in productivity was fueled by two effects: a decreased generation interval

motivated by a new technology that accurately estimates a bull's genetic traits as soon as it is born and more intense competition between animal breeders to produce high-performing bulls. Faster access to information allowed breeders to select better bulls faster, and consequently, genetic gains per generation are now much higher than they used to be in the past.

The most popular method to breed highly productive bulls is by breeding lines that descend from a famous ancestor; we call these animals supersires and show that breeding on these well-known lines is the most critical driver of a greater degree of inbreeding among bulls. As the average inbreeding rate increases, every new generation of animals experiences an increased risk of experiencing inbreeding depression, which in turn leads to lower fertility and a higher prevalence of genetic diseases. Inbreeding depression is a cost that breeders impose on others, a negative externality that is a function of the number of breeders that release closely related bulls into the market. Since pedigrees transmit the externality, it is a network externality that affects all breeders and dairy farmers.

It is impossible to directly test the impact of genomic selection on inbreeding rates because there is no natural control group and no lines unaffected by genomic selection. We used a propensity score matching method to find equivalent families based on the founder's genetic traits but had fewer offspring; breeders could have invested in them but did not. We then track those families across years and observe the average inbreeding rates of their offspring. We show that bulls from the treatment group (those who descend from famous ancestors) have consistently higher inbreeding rates than those from less-known ancestors from 2012 to 2017. Our analysis shows that the cost of this increased inbreeding amounts to around 500 million dollars.

Inbreeding rates can be reduced significantly by importing bulls from other countries that are not (or less) related to supersires. Another possible solution is to adjust the values of genetic traits by inbreeding; the Holstein Association of the United States adjusts the values of PTAs according to inbreeding levels to penalize breeders that select bulls with high inbreeding rates to account for the higher reproduction costs of such animals. The value of this penalty is a crucial element in reducing inbreeding by incentivizing dairy farmers and animal breeders to breed lines that are less related on average by, for example,

importing bulls from abroad or by crossbreeding Holsteins with other high-yield milk breeds.

8 Appendix




Figure 12: Sample selection mechanism




Figure 13: Distribution of Inbreeding rates across pre-treatment groups




Figure 14: Propensity score by class of bull

Table 1: Balance table for outcome and matched variables

Variable	Control mean (n = 224)	Treatment mean (n = 43)	t.statistic	p.value	df
Inbreeding rate	2.968	2.904	0.426	0.672	58.147
Milk	0.059	0.059	-0.036	0.971	59.956
Fat	-0.122	-0.170	0.346	0.730	76.548
Protein	-0.851	-1.009	0.416	0.679	61.059
Somatic cell score	-0.151	-0.037	-0.675	0.502	66.764
Productive life	-0.619	-0.677	0.173	0.864	59.829
Daughter preg. rate	-1.224	-0.489	-1.336	0.186	70.101
Heifer conc. rate	-11.953	-5.553	-1.807	0.075*	71.841
Cow conc. rate	-0.420	-0.209	-1.278	0.206	65.480
Livability	-0.393	-0.177	-1.816	0.074*	70.905
Type	-0.827	-0.551	-1.721	0.090*	69.510
Gest. length	0.342	-0.055	1.719	0.090*	64.087
Heifer liv.	-0.569	-0.579	0.032	0.975	58.933
Eff. calving	0.101	0.232	-1.549	0.127	58.339
Mastitis	-0.888	-0.764	-0.288	0.774	58.621
Metritis	-0.840	-0.891	0.202	0.840	62.094
Strength	-0.589	-0.581	-0.047	0.963	57.734
Rear legs rear view	-191.265	-24.936	-1.581	0.119	67.713
Rear legs side view	-0.027	-0.013	-0.495	0.623	60.964
Foot leg score	-1.348	-1.123	-0.666	0.508	59.099
Teat rear place	-8.284	-3.255	-1.859	0.067	68.421
Rump angle	-0.360	-0.136	-1.447	0.153	64.334
Thurl width	-0.010	0.050	-0.342	0.734	63.537
Foot angle	-0.767	-0.417	-2.044	0.045**	68.106
Fore udder	-0.743	-0.432	-1.735	0.087*	66.269
Rear udder height	0.142	0.193	-0.254	0.800	60.330
Rear udder width	3.044	2.995	0.690	0.493	52.692
Udder cleft	-0.386	-0.349	-0.235	0.815	77.188
Udder depth	-0.206	-0.267	0.409	0.684	67.108
Teat fron place	-0.343	-0.195	-1.006	0.318	66.497
Teat length	0.197	0.159	0.195	0.846	59.002
Milk fever	-0.312	-0.140	-0.948	0.347	62.811
Stature	-0.255	-0.233	-0.138	0.890	72.664
Dairy form	-0.573	-0.403	-1.442	0.153	73.087
Body depth	-0.401	-0.193	-1.127	0.264	61.777
Sire calv. ease	-0.738	-0.589	-0.989	0.326	73.000
Daughter calv. ease	2.427	2.287	1.188	0.239	63.963

Notes:

***Significant at the 1 percent level.

**Significant at the 5 percent level.

*Significant at the 10 percent level.

References

- Adams, H. A., Sonstegard, T. S., VanRaden, P. M., Null, D. J., Van Tassell, C. P., Larkin, D. M., and Lewin, H. A. (2016). Identification of a nonsense mutation in APAF1 that is likely causal for a decrease in reproductive efficiency in holstein dairy cattle. *Journal of Dairy Science*, 99(8):6693–6701.
- Bertrand, M., Duflo, E., and Mullainathan, S. (2004). How much should we trust differences-in-differences estimates? *The Quarterly Journal of Economics*, 119(1):249–275.
- Bourdon, R. M. (2000). *Understanding Animal Breeding*. Prentice Hall, 2 edition.
- Cameron, A. C. and Trivedi, P. (2005). *Microeconometrics: Methods and Applications*. Cambridge University Press, Cambridge, UK.
- Cunningham, S. (2021). *Causal Inference: The Mixtape*. Yale University Press, New Haven.
- Farley, J. U. (1964). "brand loyalty" and the economics of information. *The Journal of Business*, 37(4):370–381.
- Freyaldenhoven, S., Hansen, C., and Shapiro, J. M. (2019). Pre-event trends in the panel event-study design. *American Economic Review*, 109(9):3307–38.
- Guinan, F., Wiggans, G., Norman, H., Dürr, J., Cole, J., Van Tassell, C., Misztal, I., and Lourenco, D. (2023). Changes in genetic trends in us dairy cattle since the implementation of genomic selection. *Journal of Dairy Science*, 106(2):1110–1129.
- Heckman, J. J., Ichimura, H., and Todd, P. E. (1997). Matching as an econometric evaluation estimator: Evidence from evaluating a job training programme. *The Review of Economic Studies*, 64(4):605–654.
- Holstein Foundation (2018). Interpreting and Utilizing Haplotype Information. https://www.holsteinusa.com/pdf/Interpreting_and_Utilizing_Haplotype_Information_1218.pdf. Accessed: 2023-14-06.

Table 2: Difference-in-Differences coefficient estimates

	No covariates		inbreeding	
	(1)	(2)	Traits (3)	Traits and interactions (4)
treat × post = 1	1.453***			
	(0.4250)			
treat × yob = 2005		0.0944	0.0489	0.0252
		(0.2975)	(0.2989)	(0.2829)
treat × yob = 2006		-0.2183	-0.1182	-0.2789
		(0.3067)	(0.2933)	(0.2534)
treat × yob = 2007		0.2328	0.2826	0.2848
		(0.3038)	(0.2676)	(0.2534)
treat × yob = 2008		0.4035	0.3246	0.3294
		(0.3148)	(0.2798)	(0.2425)
treat × yob = 2010		-0.4589	-0.4721*	-0.5583**
		(0.2827)	(0.2826)	(0.2667)
treat × yob = 2011		-0.0164	-0.0455	-0.1671
		(0.3599)	(0.2596)	(0.2511)
treat × yob = 2012		1.224**	1.252**	1.151*
		(0.4906)	(0.5633)	(0.5981)
treat × yob = 2013		1.433***	1.083**	0.9980**
		(0.4124)	(0.4631)	(0.4725)
treat × yob = 2014		1.780***	1.600***	1.501***
		(0.3017)	(0.3514)	(0.3068)
treat × yob = 2015		1.637***	1.363***	1.255***
		(0.3453)	(0.4159)	(0.4256)
treat × yob = 2016		2.414***	1.766***	1.710***
		(0.2424)	(0.2697)	(0.2564)
treat × yob = 2017		2.849***	2.770***	2.734***
		(0.4456)	(0.4429)	(0.4507)
treat × yob = 2018		3.417***	3.576***	3.515***
		(0.4530)	(0.4566)	(0.4847)
treat × yob = 2019		2.005**	2.071**	2.022**
		(0.8611)	(0.8827)	(0.8785)
p-value for nonzero pre-effect		0.416	0.498	0.107
Observations	15,491	15,491	15,491	15,491
R ²	0.16988	0.47369	0.49942	0.50471
Within R ²	0.00702	0.01235	0.04150	0.05163
treat fixed effects	✓	✓	✓	✓
post fixed effects	✓			
year of birth fixed effects		✓	✓	✓
parent company fixed effects			✓	✓

- Holstein Foundation (2021). Understanding Genetics and the Sire Summaries. <https://forms.gle/bfB57tLEijiorH7u8>. Accessed: 2023-14-06.
- Hutchins, J. and Hueth, B. (2023). 100 years of data sovereignty: Cooperative data governance and innovation in us dairy. *Applied Economic Perspectives and Policy*, pages 1–26.
- Imbens, G. W. and Rubin, D. B. (2015). *Causal Inference for Statistics, Social, and Biomedical Sciences: An Introduction*. Cambridge University Press.
- Katz, M. L. and Shapiro, C. (1985). Network externalities, competition, and compatibility. *The American Economic Review*, 75(3):424–440.
- Kerr, W. A. (1984). Selective breeding, heritable characteristics and genetic-based technological change in the canadian beef cattle industry. *Western Journal of Agricultural Economics*, 9(1):14–28.
- Ladd, G. W. and Martin, M. B. (1976). Prices and demands for input characteristics. *American Journal of Agricultural Economics*, 58(1):21–30.
- Lewbel, A. (2019). The identification zoo: Meanings of identification in econometrics. *Journal of Economic Literature*, 57(4):835–903.
- Liebowitz, S. J. and Margolis, S. E. (1994). Network externality: An uncommon tragedy. *The Journal of Economic Perspectives*, 8(2):133–150.
- Lush, J. L. (1937). *Animal Breeding Plans*. Collegiate Press, Incorporated.
- Melton, B. E., Colette, W. A., and Willham, R. L. (1994). Imputing input characteristic values from optimal commercial breed or variety choice decisions. *American Journal of Agricultural Economics*, 76(3):478–491.
- Melton, B. E., Heady, E. O., and Willham, R. L. (1979). Estimation of economic values for selection indices. *Animal Production*, 28(3):279–286.
- Montiel-Olea, J. L. and Plagborg-Møller, M. (2019). Simultaneous confidence bands: Theory, implementation, and an application to SVARs. *Journal of Applied Econometrics*, 34(1):1–17.

- Richards, T. J. and Jeffrey, S. R. (1996). Establishing Indices of Genetic Merit Using Hedonic Pricing: An Application to Dairy Bulls in Alberta. *Canadian Journal of Agricultural Economics/Revue canadienne d'agroeconomie*, 44(3):251–264.
- Rosen, S., Murphy, K. M., and Scheinkman, J. A. (1994). Cattle cycles. *Journal of Political Economy*, 102(3):468–492.
- Rosenbaum, P. R. and Rubin, D. B. (1983). The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70(1):41–55.
- Roth, J. (2022). Pretest with caution: Event-study estimates after testing for parallel trends. *American Economic Review: Insights*, 4(3):305–22.
- Roth, J., Sant'Anna, P. H., Bilinski, A., and Poe, J. (2023). What's trending in difference-in-differences? a synthesis of the recent econometrics literature. *Journal of Econometrics*, 235(2):2218 – 2244.
- Schroeder, T. C., Espinosa, J. A., and Goodwin, B. K. (1992). The value of genetic traits in purebred dairy bull services. *Review of Agricultural Economics*, 14(2):215–226.
- Smith, J. A. and Todd, P. E. (2005). Does matching overcome Lalonde's critique of non-experimental estimators? *Journal of Econometrics*, 125(1):305–353. Experimental and non-experimental evaluation of economic policy and models.
- Steyn, Y., Lawlor, T. J., Lourenco, D., and Misztal, I. (2023). The importance of historically popular sires on the accuracy of genomic predictions of young animals in the US Holstein population. *Journal of Dairy Science Communications*, 4(4):260–264.
- Steyn, Y., Masuda, Y., Tsuruta, S., Lourenco, D., Misztal, I., and Lawlor, T. (2022). Identifying influential sires and distinct clusters of selection candidates based on genomic relationships to reduce inbreeding in the us holstein. *Journal of Dairy Science*, 105(12):9810–9821.
- Stigler, G. J. (1961). The economics of information. *Journal of Political Economy*, 69(3):213–225.

Sun, L. and Abraham, S. (2021). Estimating dynamic treatment effects in event studies with heterogeneous treatment effects. *Journal of Econometrics*, 225(2):175 – 199.

Sy, H. A., Faminow, M. D., Johnson, G. V., and Crow, G. (1997). Estimating the Values of Cattle Characteristics using an Ordered Probit model. *American Journal of Agricultural Economics*, 79(2):463–476.

Wiggans, G. R. and Carrillo, J. A. (2022). Genomic selection in United States dairy cattle. *Frontiers in Genetics*, 13.